

Drugs of Abuse Table URINE DRUG TESTING

DRUGCHECK[®]

Drug Type/Street Names	Brand Name	Detection Time	Signs of Use
Alcohol <small>.C ALC ALC ALC</small>			
Booze, Brew, Cold One, Juice, Sauce, Vino, Hard Stuff	—	< 1 day	Odor of alcohol on breath or in sweat, slurred speech, lack of coordination, poor balance, red eyes or flushed face, nausea, weakness, or sweatiness
Amphetamine <small>AP AMP AMP AMP</small>			
Speed, Bennies, Dexies, Uppers, Crank, Ice, Black Beauties	Dexedrine, Benzedrine, Dextrostat, Adderall, Biphemamine, Ritalin	Single use: 1-2 days Prolonged use: 7-10 days	Increased blood pressure and pulse rate, insomnia, loss of appetite and physical exhaustion
Barbiturates <small>IR BAR BAR BAR BAF</small>			
Barbs, Downers, Yellow Jackets, Reds, Red Devils, Goof Balls, Reds & Blues, Peanuts	Nembutal, Seconal, Butisol, Florinal, Phenobarbital	2 days (Nembutal, Seconal, Butisol, Florinal) 1-3 weeks (Phenobarbital)	Mild euphoria, lack of inhibition, relief of anxiety and sleepiness. Higher doses cause impairment of memory, judgement and coordination, irritability, and paranoid and suicidal ideation
Benzodiazepines <small>ZO BZO BZO BZO</small>			
Benzos, Downers, Tranks, Blues, Yellows	Valium, Serax, Xanax, Klonopin, Librium, Ativan, Loraz, Halcion, Rohypnol, Restoril, Versed, Rivotril, Dalamane, Halcion	5-7 days 5-14 days (Klonopin) 3 days (Rohypnol)	Amnesia, hostility, irritability, sleepiness and vivid or disturbing dreams
Buprenorphine <small>JP BUP BUP BUP</small>			
—	Suboxone, Subutex, Buprenex	2-4 days	Pinpoint pupils, extreme drowsiness, dizziness, blurred vision, slowed breathing
Cocaine <small>OC COC COC COC</small>			
Coke, Crack, Snow, Flake, Rock, Powder, Blow, Soda Cot	—	Single use: 3 days Prolonged use: 4 days	Increased blood pressure and heart rate, dilated pupils, insomnia, loss of appetite, cardiac arrhythmias, ischemic heart conditions, sudden cardiac arrest, convulsions and stroke
Cotinine <small>JT COT COT COT</small>			
(Metabolite of nicotine)	—	2-6 days ¹	—
EDDP <small>IDP EDDP EDDP EDC</small>			
(Metabolite of methadone)	—	Up to 7 days ²	EDDP is the primary metabolite of methadone; symptoms of methadone use: Restlessness, vomiting, nausea, slowed breathing, itchy skin, pupil contraction, severe sweating ³
Ethyl Glucuronide <small>G ETG ETG ETG</small>			
—	—	Up to 80 hours	EtG is a unique biomarker of alcohol use (see "Alcohol" above)
Fentanyl <small>L FYL FYL FYL</small>			
Dance Fever, Friend, Tango and Cash, China Girl, Goodfella, Jackpot, TNT	Actiq, Duragesic, Sublimaze	1-2 days ⁴	Drowsiness, dizziness, lightheadedness, false sense of well being ⁵
Gamma Hydroxybutyric Acid <small>B GHB GHB GHB GHI GHB GHB GHB GHB C</small>			
GHB, Easy Lay, Georgia Home Boy, Grievous Bodily Harm, Scoop, Goop, Fantasy, Everclear, Liquid X	—	12 hours	Euphoria, drowsiness, decreased anxiety, confusion, memory impairment, visual hallucinations and paradoxically and excited and aggressive behavior
K2/Spice – Synthetic Cannabis <small>C2/SPICE K2/SPICE CE K2/SPICE K2/E</small>			
Bliss, Black Mamba, Bombay Blue, Fake Weed, Genie, Spice, Zohai	—	Up to 3 days	Increased heart rate and blood pressure, paranoia, panic attacks and giddiness
Ketamine <small>ET KET KET KET</small>			
Kat, Special K, Vitamin K, Cat Valium, Kit Kat, K	Ketalar	< 3 days	Hallucinations, involuntary rapid eye movement, dilated pupils, salivation, tear secretions and stiffening of the muscles
Marijuana <small>IC THC THC THC</small>			
Weed, Pot, Grass, Reefer, Hash, Mary Jane, Cannabis, Colombian, Acapulco Gold, Ganja, Bhang	Marinol	Single use: 2-7 days Prolonged use: 1-2 months	Increased appetite, altered perception, memory loss, bloodshot eyes, giggle frequently, trouble walking, disorientation; increased heart rate, vasodilation, dry mouth and throat, decreased respiratory rate, panic attacks, and paranoia

Drug Type/Street Names	Brand Name	Detection Time	Signs of Use
Methadone ^{MTD} ^{MTD} ^{MTD} ^{MTD} ^{MTD}			
Amidone, Fizzes, Maria, Wafer	Dolophine, Methadose, Amidone, L-Polamidon, Physeptone	3 days	Used to treat opioid addiction and pain; significant overdose risk when used improperly. Sweating, itchy skin or sleepiness
Methamphetamine ^{MET} ^{MET} ^{MET} ^{MET}			
Speed, Meth, Crystal Ice, Crank, Tweak, Glass, Poor Man's Cocaine	Desoxyn, Methedrine	Single use: 1-2 days Prolonged use: 7-10 days	Increased wakefulness, increased physical activity, decreased appetite, rapid breathing and heart rate, irregular heartbeat, increased blood pressure and hyperthermia (overheating)
Methylenedioxyamphetamine (Ecstasy) ^{MDMA} ^{MDMA} ^{MDMA} ^{MDMA}			
XTC, E, Adam, Disco Biscuit, Hug Drug, Lover's Speed, Beans, Clarity, Eve	None	1 day	Same as amphetamines: euphoria, increased sensitivity to touch, sensual and sexual arousal, confusion, anxiety, depression, paranoia, involuntary teeth clenching, tremors, chills, sweating and blurred vision
Opiates ^{OPI} ^{OPI} ^{OPI} ^{OPI}			
(Heroin- and morphine-based drugs) Smack, H, Skag, Dope, Junk, Horse	Hydromorphone (Dilaudid), Codeine, Tylenol with Codeine, Morphine, Avinza, Kadian, MS Contin, Oramorph SR, Hydrocodone (Vocodin): Pos. at 300 cutoff	1-2 days	Constipation, pupillary constriction, urinary retention, nausea, vomiting, respiratory depression, dizziness, impaired coordination, loss of appetite, rash, slow or rapid heartbeat and changes in blood pressure
Oxycodone ^{OXY} ^{OXY} ^{OXY} ^{OXY}			
OXY, OC, OX, Oxycotton, Hillbilly Heroin, Killer	OxyContin, Percodan, Percocet, Roxicodone, Tylox, OxyContin is a time-released form of Oxycodone	1-3 days	Respiratory depression, constipation, papillary constriction and cough suppression. Extended chronic use of oxycodone containing acetaminophen may cause severe liver damage
Phencyclidine ^{PCP} ^{PCP} ^{PCP} ^{PCP}			
Veterinary Anesthetic, Angel Dust, Hog, Killer Weed	Sernylan	Single use: 1 week Prolonged use: 2-4 weeks	Elevated blood pressure, sweating, involuntary rapid eye movements of the eyeball, nausea and vomiting, extreme aggressive behavior, euphoria, intoxicated appearance
Propoxyphene ^{PPX} ^{PPX} ^{PPX} ^{PPX}			
Yellow, Footballs	Darvon, Darvocet	6 hours to 2 days	Blurred vision, headache, nausea and vomiting, euphoria, intoxication, mild depression, seizures, confusion, hallucination and delusions
Tramadol ^{TML} ^{TML} ^{TML} ^{TML}			
–	ConZip, Tramal, Ultram, Ultram ER	Undocumented	Nausea, vomiting, drowsiness, dry mouth, perspiration, dizziness, tremor, confusion, hallucination, blood pressure instability, and exacerbation of convulsions ⁶
Tricyclic Antidepressants ^{TCA} ^{TCA} ^{TCA} ^{TCA}			
–	Amitriptyline (Elavil, Endep), Cyclobenzaprine (Flexeril), Clomipramine (Anafranil), Perphenazine (Triafron), Desipramine (Norpramin), Imipramine (Tofranil), Nortriptyline (Pamelor), Protriptyline (Vivactil), Trimipramine (Surmontil), Promazine (Sparine), Nordoxepine, Doxepin (Sinequan)	2-7 days ⁷	Dry mouth, blurred vision, drowsiness, lightheadedness (drop in blood pressure when moving from sitting to standing), increased sweating ⁸

REFERENCES

1. University of Minnesota Medical Center, "Nicotine-Cotinine," 2015. Web. 1 June 2015.
2. Mayo Medical Laboratories, "Drug Book: Methadone," 2015. Web. 1 June 2015.
3. University of Maryland Center for Substance Abuse Research, "Drugs: Methadone," 2013. Web. 1 June 2015.
4. Silverstein, J.H. et. al., "An analysis of the duration of fentanyl and its metabolites in urine and saliva," 1993. Web. 1 June 2015.
5. Mayo Clinic, "Drugs-Supplements: Fentanyl," 2015. Web. 1 June 2015.

6. Rajabizadeh, Ghodratalah, M.D. et. al., 2009, National Institutes of Health. Web. 1 June 2015.
7. Ningegowda, Lokesh, M.D., Cleveland Clinic. Web. 1 June 2015.
8. Mayo Clinic, "Diseases-Conditions: Depression: Antidepressants," 2015. Web. 1 June 2015.



**ROYAL
MEDICAL**

"Improving healthcare and workplace safety through innovation".

Australian Importer and Provider of Technical Support

T: (+612) 9939 4122
F: (+612) 9939 4133
E: info@royalmedical.com.au

Contact us today:

Our websites:
www.royalmedical.com.au
www.royalmedical.co.nz
www.drugcheck.com.au
www.drug-test.com.au